

GOSAT 搭載フーリエ変換分光計による大気中温室効果ガスの観測
(国立環境研究所) ○森野勇・井上誠・中前久美・菊地信弘・吉田幸生・内野修・
横田達也

Observations of atmospheric greenhouse gases with a Fourier transform spectrometer onboard GOSAT (Greenhouse Gases Observing SATellite) (National Institute for Environmental Studies) Isamu Morino, Makoto Inoue, Kumi Nakamae, Nobuhiro Kikuchi, Yukio Yoshida, Osamu Uchino, Tatsuya Yokota

The Greenhouse gases Observing SATellite (GOSAT), launched on 23 January 2009, is the world's first satellite dedicated to measuring the concentrations of the two major greenhouse gases, carbon dioxide (CO_2) and methane (CH_4), from space. The data measured with the Thermal And Near-infrared Sensor for carbon Observation Fourier Transform Spectrometer (TANSO-FTS) and the Cloud and Aerosol Imager (TANSO-CAI) are processed into several types of data products. Column abundances of CO_2 and CH_4 , Level 2 FTS SWIR (Short-Wavelength InfraRed) data product are retrieved from the Level 1B FTS SWIR spectral data. Validation of the GOSAT TANSO-FTS SWIR Level 2 products is critical since the data are used for generating the Level 3 FTS product, global distributions of column-averaged mixing ratio of CO_2 and CH_4 ($X\text{CO}_2$ and $X\text{CH}_4$), and the Level 4 product, regional CO_2 fluxes and three dimensional distribution of CO_2 calculated from the estimated fluxes. Yokota et al. [1] reported the first preliminary result observed with GOSAT TANSO-FTS SWIR.

Preliminary validation of the GOSAT TANSO-FTS SWIR Level 2 products was made with ground-based high-resolution FTS data of the Total Carbon Column Observing Network (TCCON) data [2]. $X\text{CO}_2$ and $X\text{CH}_4$ retrieved with the TANSO-FTS SWIR ver. 01.xx retrieval algorithm showed large negative biases and standard deviations (-8.85 ppm and 4.75 ppm for $X\text{CO}_2$ and -20.4 ppb and 18.9 ppb for $X\text{CH}_4$, respectively) compared with the TCCON data [3].

Therefore we investigated the influence of aerosols and thin cirrus clouds on $X\text{CO}_2$ by lidar and sky radiometer at Tsukuba TCCON site (36.051N, 140.122E), and found that it was important to take into account vertical profiles of aerosols and thin cirrus clouds and to use more adequate solar irradiance database in order to improve the GOSAT $X\text{CO}_2$ data [4].

Based on these results and sensitivity analysis of the retrieval algorithm, multiple reasons for these error characteristics (e.g., solar irradiance database, handling of aerosol scattering) are identified and corrected in a revised version of the TANSO-FTS SWIR retrieval algorithm (ver. 02.xx). The improved retrieval algorithm shows much smaller biases and standard deviations (-1.48 ppm and 2.09 ppm for $X\text{CO}_2$ and -5.9 ppb and 12.6 ppb for $X\text{CH}_4$, respectively) than those of the ver. 01.xx. [5].

To improve further the retrieval algorithm, we have started to investigate the impact of aerosols and thin cirrus clouds on the GOSAT data over Tsukuba and Saga

(33.241N, 130.288E) in Japan and Lauder (45.038S, 169.684E) in New Zealand by high-resolution ground-based FTS, lidar and sky radiometer.

We will present the latest results on the GOSAT observations and activities of validating the GOSAT TANSO-FTS SWIR Level 2 products with TCCON and aircraft data and improving the retrieval algorithm.

References

- [1] T. Yokota, Y. Yoshida, N. Eguchi, Y. Ota, T. Tanaka, H. Watanabe, and S. Maksyutov, *SOLA*, **5**, 160–163, doi:10.2151/sola.2009-041, 2009.
- [2] D. Wunch, G. C. Toon, J.-F. L. Blavier, R. A. Washenfelder, J. Notholt, B. J. Connor, D. W. T. Griffith, V. Sherlock, and P. O. Wennberg, *Philos. T. Roy. Soc. A*, **369**, 2087–2112, doi:10.1098/rsta.2010.0240, 2011.
- [3] I. Morino, O. Uchino, M. Inoue, Y. Yoshida, T. Yokota, P. O. Wennberg, G. C. Toon, D. Wunch, C. M. Roehl, J. Notholt, T. Warneke, J. Messerschmidt, D. W. T. Griffith, N. M. Deutscher, V. Sherlock, B. Connor, J. Robinson, R. Sussmann, and M. Rettinger, *Atmos. Meas. Tech.* **4**, 1061–1076, doi:10.5194/amt-4-1061-2011, 2011.
- [4] O. Uchino, N. Kikuchi, T. Sakai, I. Morino, Y. Yoshida, T. Nagai, A. Shimizu, T. Shibata, A. Yamazaki, A. Uchiyama, N. Kikuchi, S. Oshchepkov, A. Bril, and T. Yokota, *Atmos. Chem. Phys.*, **12**, 3393–3404, doi:10.5194/acp-12-3393-2012, 2012.
- [5] Y. Yoshida, N. Kikuchi, I. Morino, O. Uchino, S. Oshchepkov, A. Bril, T. Saeki, N. Schutgens, G. C. Toon, D. Wunch, C. M. Roehl, P. O. Wennberg, D. W. T. Griffith, N. M. Deutscher, T. Warneke, J. Notholt, J. Robinson, V. Sherlock, B. Connor, M. Rettinger, R. Sussmann, P. Ahonen, P. Heikkinen, E. Kyrö, and T. Yokota, *Atmos. Meas. Tech. Discuss.*, **6**, 949–988, doi:10.5194/amtd-6-949-2013, 2013.